arXiv:1211.1727v1 [math.NT] 7 Nov 2012

AN ALTERNATIVE PROOF AND GENERALIZATION OF FERRERO'S COMPUTATIONS OF IWASAWA λ-INVARIANTS

JORDAN SCHETTLER

Abstract. We prove a slight generalization of Iwasawa's 'Riemann-Hurwitz' formula for number fields and use it to generalize Ferrero's computations of Iwasawa λ-invariants for the cyclotomic $\mathbb{Z}_2$-extensions of imaginary quadratic number fields. In particular, we show that if p is a Fermat prime, then similar computations of Iwasawa λ-invariants hold for certain imaginary quadratic extensions of the unique subfield $k \subset \mathbb{Q}(\zeta_{p^2})$ such that $[k : \mathbb{Q}] = p$.

1. Introduction

Fix a prime p and number field k. Suppose k_∞ is a $\mathbb{Z}_p$-extension of k, i.e., k_∞ is a Galois extension of k with Galois group $\mathrm{Gal}(k_\infty/k)$ isomorphic to the group $\mathbb{Z}_p$ of p-adic integers. The subfields of k_∞ which contain k all lie in a tower

$$k = k_0 \subset k_1 \subset k_2 \subset \ldots \subset k_\infty$$

with $\mathrm{Gal}(k_n/k) \cong \mathbb{Z}/(p^n)$ for all integers $n \geq 0$. Kenkichi Iwasawa's well-known growth formula (see [Iwa59] or [Iwa73a]) states that there are integers λ, μ, ν such that if A_n denotes the p-primary part of the class group of k_n, then

$$|A_n| = p^{\lambda \cdot n + \mu \cdot p^n + \nu}$$

for all sufficiently large integers n. In particular, $\lambda, \mu \geq 0$ but we can have $\nu < 0$. We call λ, μ, ν the Iwasawa invariants of the extension k_∞/k. There is a special case in which all the invariants are known to vanish.

Theorem 1 (Iwasawa, [Iwa59]). *Let k be a number field having exactly one prime $\mathfrak{p}$ lying over a rational prime p, and let k_∞ be a $\mathbb{Z}_p$-extension of k. Suppose p does not divide the class number of k. Then for every number field $k_n \subset k_\infty$ which contains k, the prime $\mathfrak{p}$ ramifies totally in k_n/k and p does not divide the class number of k_n; in particular, all of the Iwasawa invariants for k_∞/k are zero, i.e., $\lambda = \mu = \nu = 0$.*

For every prime p and number field k, there is at least one $\mathbb{Z}_p$-extension of k; namely, there is a unique $\mathbb{Z}_p$-extension of k contained in $\bigcup_{n\geq 0} k(\zeta_{p^n})$ where the ζ_{p^n} are primitive p^nth roots of unity. We denote this $\mathbb{Z}_p$-extension by k_∞^{cyc} and call it the cyclotomic $\mathbb{Z}_p$-extension of k. We write $\lambda_p(k)$, $\mu_p(k)$, $\nu_p(k)$ for the Iwasawa invariants of the extension $k_\infty^{\mathrm{cyc}}/k$.

Iwasawa conjectured that $\mu_p(k) = 0$ for all primes p and number fields k, and no counterexamples are known.[1] This conjecture has been verified for abelian number fields by Bruce Ferrero and Lawrence Washington ([FW79]) and for p-extensions of number fields k with $\mu_p(k) = 0$ by Iwasawa ([Iwa73b]). We make the assumption $\mu_p(k) = 0$ throughout the paper.

---

[1] However, there are non-cyclotomic $\mathbb{Z}_p$-extensions of number fields which have an Iwasawa invariant $\mu > 0$.

Bruce Ferrero and Yûji Kida independently calculated $\lambda_2(k)$ for imaginary quadratic fields k. Their computations are explicit:

Theorem 2 (Ferrero, [Fer80]; Kida, [Kid79])**.** *Let $d > 2$ be a squarefree integer. Then*

$$\lambda_2(\mathbb{Q}(\sqrt{-d})) = -1 + \sum_{\substack{p|d \\ p\neq 2}} 2^{\mathrm{ord}_2(p^2-1)-3}$$

where the sum ranges over all odd primes p dividing d.

Here $\mathrm{ord}_2(n)$ denotes the 2-adic order of an integer n, i.e., the largest exponent e such that $2^e|n$. Thus, for example, if p is a prime with $p \equiv \pm 3 \pmod 8$, then $\mathrm{ord}_2(p^2 - 1) = 3$, so $\lambda_2(\mathbb{Q}(\sqrt{-p})) = -1 + 2^{3-3} = 0$. In fact, there are infinitely many such primes, so for any integer $m \geq 0$ there are infinitely many choices of $m+1$ such distinct primes $p_1, \ldots, p_{m+1}$, and we get $\lambda_2(\mathbb{Q}(\sqrt{-p_1 p_2 \cdots p_{m+1}})) = m$.

Iwasawa argued in [Iwa65] that when $\mu_p(k) = 0$, the invariant $\lambda_p(k)$ is a good analog for twice the genus of a curve; we expound on this idea more precisely below. In fact, Iwasawa derived a 'Riemann-Hurwitz' formula for cyclic, degree p extensions of number fields in which the invariant $\lambda_p(k)$ does indeed correspond to twice the genus of a curve in the classical Riemann-Hurwitz formula. Below we prove a slight generalization of Iwasawa's 'Riemann-Hurwitz' formula, and then use the formula to generalize Theorem 2 of Ferrero and Kida.

2. Iwasawa's 'Riemann-Hurwitz' Formula Revisited

Following Iwasawa, we define a $\mathbb{Z}_p$-field to be the cyclotomic $\mathbb{Z}_p$-extension field of a number field. In other words, K is a $\mathbb{Z}_p$-field when $K = k_\infty^{\mathrm{cyc}}$ for the prime p. Note that if $k_\infty^{\mathrm{cyc}} = \ell_\infty^{\mathrm{cyc}}$ for some prime p and number fields k, ℓ, then $\lambda_p(k) = \lambda_p(\ell)$ and we have $\mu_p(k) = 0 \Leftrightarrow \mu_p(\ell) = 0$. Thus for a $\mathbb{Z}_p$-field K we may define λ_K to the be Iwasawa λ-invariant $\lambda_p(k)$ for any number field k with $K = k_\infty^{\mathrm{cyc}}$. Likewise, we may define the notation $\mu_K = 0$ to indicate that $\mu_p(k) = 0$ for some (and hence every) number field k such that $K = k_\infty^{\mathrm{cyc}}$.

Theorem 3. *Suppose K is a $\mathbb{Z}_p$-field. Let I_K denote the group of invertible fractional ideals in the integer ring $\mathcal{O}_K$ of K, and let $P_K \leq I_K$ denote the subgroup of principal fractional ideals. Then the p-primary part A_K of the class group $C_K = I_K/P_K$ satisfies*

$$A_K \cong (\mathbb{Q}_p/\mathbb{Z}_p)^{\lambda_K} \oplus M$$

where $p^n M = 0$ for some $n \geq 0$. Moreover, M is trivial precisely when $\mu_K = 0$.

This structure theorem, whose proof may be gleaned from [Iwa59] and [Iwa73a], allows us to pin down the analogy between number fields and curves in the spirit of [Iwa65]. The analogy is illustrated in Table 1. A few remarks are in order. The ring $\mathcal{O}_K$ is not Noetherian, but by inverting p, the resulting ring $\mathcal{O}_K[1/p]$ is actually a Dedekind domain, so its prime spectrum is a nice scheme which shares many properties with curves. This also helps to explain why the ramification for the prime p is missing in Iwasawa's 'Riemann-Hurwitz' formula; however, we will not use this interpretation in the proof of Iwasawa's formula, but the so-called Dedekind

If K is the function field of a smooth, projective curve X over $\mathbb{C}$, then the local rings of X at closed points are DVRs, and the p-primary part of the Picard group satisfies $$\mathrm{Pic}(X)[p^\infty] \cong (\mathbb{Q}_p/\mathbb{Z}_p)^{2g_K}$$ where g_K is the genus.	If K is a $\mathbb{Z}_p$-field with $\mu_K = 0$ and $X = \mathrm{Spec}(\mathcal{O}_K[1/p])$, then the local rings of X at closed points are DVRs, and the p-primary part of the Picard group satisfies $$\mathrm{Pic}(X)[p^\infty] \cong (\mathbb{Q}_p/\mathbb{Z}_p)^{\lambda_K}$$ where λ_K is an Iwasawa invariant.

Table 1. Similarities in structures of Picard groups

different formula implies that a purely geometric proof of Iwasawa's result should exist. Before stating Iwasawa's 'Riemann-Hurwitz' formula, we need a definition.

Definition 4. Let G be a cyclic p-group. For a G-module M, define the 'Euler characteristic' $\chi(G, M) \in \mathbb{Z}$ to be the exponent of p in the Herbrand quotient

$$p^{\chi(G,M)} = \frac{|H^2(G, M)|}{|H^1(G, M)|}$$

when both cohomology groups $H^i(G, M)$ are finite for $i = 1, 2$. Note that χ inherits the following properties from the Herbrand quotient:

1. χ is additive on short exact sequences of G-modules[2]
2. $\chi(G, M) = 0$ when M is a finite G-module
3. $\chi(G, M^*) = -\chi(G, M)$ when $M^* = \mathrm{Hom}_{\mathbb{Z}_p}(M, \mathbb{Q}_p/\mathbb{Z}_p)$ is the p-Pontryagin dual of a $\mathbb{Z}_pG$-module M.

Theorem 5 (Iwasawa's 'Riemann-Hurwitz' Formula)**.** *Let L/K be a $\mathbb{Z}/(p)$-extension of $\mathbb{Z}_p$-fields with $G = \mathrm{Gal}(L/K)$. Suppose $\mu_K = 0$. Then $\mu_L = 0$ and*

$$\lambda_L = p\lambda_K - (p-1)\chi(G, P_L) + \sum_{w \nmid p}(e(w) - 1) \tag{5.1}$$

where the sum ranges over all finite places w of L not lying above p, $e(w)$ is the ramification index of w in L/K, and G acts in the obvious way on the principal fractional ideals P_L of the integer ring $\mathcal{O}_L$ in L.

Remark 6. If the extension L/K in Theorem 5 is unramified at the infinite places (always true, e.g., when p is odd), then Iwasawa showed that $H^2(G, L^\times) = 0$, so additivity of the 'Euler characteristic' χ shows that $-\chi(G, P_L) = \chi(G, \mathcal{O}_L^\times)$, which recovers the formula as originally stated by Iwasawa in [Iwa81].

Proof of Theorem 5. Using the notation of Theorem 3 above, we let A_L denote the p-primary part of the class group $C_L = I_L/P_L$ where again I_L, P_L are the groups of invertible and principal, respectively, fractional ideals of the integer ring $\mathcal{O}_L$. The statement that $\mu_L = 0$ follows from $\mu_K = 0$ by a result mentioned above (see [Iwa73b]) since L/K is a p-extension, so the structure theorem implies

$$A_L^* = \mathrm{Hom}_{\mathbb{Z}_p}(A_L, \mathbb{Q}_p/\mathbb{Z}_p) \cong \mathrm{Hom}_{\mathbb{Z}_p}((\mathbb{Q}_p/\mathbb{Z}_p)^{\lambda_L}, \mathbb{Q}_p/\mathbb{Z}_p) \cong \mathbb{Z}_p^{\lambda_L}$$

---

[2] If $0 \to A \to B \to C \to 0$ is a SES of G-modules and two of the numbers $\chi(G, A)$, $\chi(G, B)$, $\chi(G, C)$ are finite (i.e., well-defined), then so is the other and $\chi(G, A) - \chi(G, B) + \chi(G, C) = 0$.

	$\mathrm{rank}_{\mathbb{Z}_p}(-)$	$(-)^G$	$H^2(G,-)$	$H^1(G,-)$	$\chi(G,-)$
$\mathbb{Z}_p$	1	$\mathbb{Z}_p$	$\mathbb{Z}_p/p\mathbb{Z}_p$	0	1
$\mathbb{Z}_pG$	p	$\mathbb{Z}_p$	0	0	0
I_pG	$p-1$	0	0	$\mathbb{Z}_p/p\mathbb{Z}_p$	-1

Table 2.

as $\mathbb{Z}_p$-modules. On the other hand, we have the following classification theorem.

Theorem 7 (Diederichsen, [Die40]). *Let $\langle g\rangle = G \cong \mathbb{Z}/(p)$. The only indecomposable $\mathbb{Z}_pG$-modules which are free of finite rank over $\mathbb{Z}_p$ are (up to isomorphism) $\mathbb{Z}_p$, $\mathbb{Z}_pG$, and the augmentation ideal $I_pG = (g-1)\mathbb{Z}_pG$.*

Hence

$$A_L^* \cong \mathbb{Z}_p^a \oplus (\mathbb{Z}_pG)^b \oplus (I_pG)^c$$

as $\mathbb{Z}_pG$-modules for some nonnegative integers a, b, c. It is easy to compute the $\mathbb{Z}_p$-ranks, G-invariants, and Euler characteristics, of these indecomposables. The results are summarized in Table 2. Moreover, if S is the set of finite places of K which do not lie above p and ramify in L/K, then Iwasawa showed in [Iwa81] that

$$\chi(G, I_L) = |S|.$$

Also, the last column in Table 2 implies

$$\chi(G, A_L^*) = a\cdot 1 + b\cdot 0 + c\cdot(-1) = a - c.$$

Hence duality gives

$$\chi(G, A_L) = -\chi(G, A_L^*) = -a + c,$$

but

$$\chi(G, C_L) = \chi(G, A_L)$$

since G is a p-group and C_L is torsion, so $\chi(G, P_L)$ is also finite and additivity gives

$$-\chi(G, P_L) + |S| = -\chi(G, P_L) + \chi(G, I_L) = \chi(G, C_L) = -a + c.$$

Also, the natural map

$$A_K \to A_L^G$$

has finite kernel and finite cokernel since the same is true of $C_K \to C_L^G$ as seen from the snake lemma. Thus the first two columns in Table 2 show that

$$\begin{aligned}\lambda_K &= \mathrm{rank}_{\mathbb{Z}_p}(A_K^*) = \mathrm{rank}_{\mathbb{Z}_p}((A_L^G)^*) = \mathrm{rank}_{\mathbb{Z}_p}((A_L^*)^G)\\ &= a\cdot 1 + b\cdot 1 + c\cdot 0 = a + b.\end{aligned}$$

Putting all of this together, we get

$$\begin{aligned}\lambda_L &= a\cdot 1 + b\cdot p + c(p-1) = p(a+b) + (p-1)(-a+c)\\ &= p\lambda_K - (p-1)\chi(G, P_L) + (p-1)|S|,\end{aligned}$$

as needed. We have shown more than just a formula for the λ-invariants; in fact,

$$A_L^* \cong \mathbb{Z}_p^a \oplus (\mathbb{Z}_pG)^{\lambda_K - a} \oplus (I_pG)^{|S| - \chi(G, P_L) + a}$$

as $\mathbb{Z}_pG$-modules for some nonnegative integer a with $\chi(G, P_L) - |S| \le a \le \lambda_K$. $\square$

3. Main Result

Theorem 8. *Suppose* p *is a prime of the form* $2^t + 1$ *for some integer* $t \geq 0$ *and let* $d > 2 \geq (d, p)$ *be a squarefree integer. Take* k *to be the unique real subfield of* $\mathbb{Q}(\zeta_{2p^2})$ *such that* $[k : \mathbb{Q}] = p$*, and take* $K = k_\infty^{\mathrm{cyc}}$ *to be the cyclotomic* $\mathbb{Z}_2$*-extension of* k*. If the class number* h_k *of* k *is odd (e.g., we can take* $p \in \{2, 3, 5, 17, 257\}$*), then*

$$\lambda_2(k(\sqrt{-d})) = -1 + |S|$$

where S *is the set of finite places of* K *not lying above* 2 *which ramify in* $K(\sqrt{-d})/K$*.*

Remark 9. Note that the field k in Theorem 8 is the first layer in the cyclotomic $\mathbb{Z}_p$-extension of $\mathbb{Q}$. Thus when $p = 2$, Theorem 8 precisely recovers Theorem 2 since, in that case, $K = k_\infty^{\mathrm{cyc}} = \mathbb{Q}_\infty^{\mathrm{cyc}}$ is the cyclotomic $\mathbb{Z}_2$-extension of $\mathbb{Q}$ and for each odd prime q there are exactly $2^{\mathrm{ord}_2(q^2-1)-3}$ primes of $\mathbb{Q}_\infty^{\mathrm{cyc}}$ which lie above q.

Proof of Theorem 8. We apply Iwasawa's 'Riemann-Hurwtiz' formula (Equation 5.1) to the extension L/K of $\mathbb{Z}_2$-fields where we take $L = K(\sqrt{-d}) = \ell_\infty^{\mathrm{cyc}}$ to be the cyclotomic $\mathbb{Z}_2$-extension of $\ell = k(\sqrt{-d})$ and as above $G = \mathrm{Gal}(L/K)$. We get

$$\lambda_2(k(\sqrt{-d})) = \lambda_L = 2\lambda_K - \chi(G, P_L) + |S|.$$

Thus it suffices to show that $\lambda_K = 0$ and $\chi(G, P_L) = 1$.

First, we prove that $\lambda_K = 0$ using Theorem 1. We have assumed that h_k is odd, and, in fact, this assumption is valid[3] for $p \in \{2, 3, 5, 17, 257\}$. Hence it is enough to show that k has exactly one prime lying above 2. When $p = 2$, this is clear since $k = \mathbb{Q}(\sqrt{2})$, so we may assume $p = 2^t + 1$ for some integer $t \geq 1$. Of course, we must have $t = 2^r$ (i.e, $p = F_r = 2^{2^r} + 1$ is a prime Fermat number) for some integer $r \geq 0$. For every $j \geq 1$, we can factor

$$F_j - 2 = 2^{2^j} - 1 = F_0 F_1 \cdots F_{j-1}$$

as a product of consecutive Fermat numbers $F_i = 2^{2^i} + 1$. This identity shows that Fermat numbers are pairwise relatively prime, so $p^2 \nmid 2^{2^j} - 1$ for any $j \geq 0$ since $p = F_r$ is a prime Fermat number. This means that the multiplicative order of 2 modulo p^2 is not a power of 2 which forces the residue degree of 2 in $\mathbb{Q}(\zeta_{p^2}) = \mathbb{Q}(\zeta_{2p^2})$ to be divisible by p. Consequently, the residue degree of 2 in k is p, which is equivalent to 2 being inert in k. Hence $\lambda_K = 0$ and it remains only to show that $\chi(G, P_L) = 1$.

To prove $\chi(G, P_L) = 1$, we will show that $|H^1(G, P_L)| = 1$ and $|H^2(G, P_L)| = 2$. First, we fix some notation. Write out the towers for the $\mathbb{Z}_2$-extensions K/k and $L/\ell = K(\sqrt{-d})/k(\sqrt{-d})$ as

$$k \subset k_1 \subset k_2 \subset \ldots \subset k_\infty := K$$
$$\ell \subset \ell_1 \subset \ell_2 \subset \ldots \subset \ell_\infty := L.$$

In this way,

$$\mathrm{Gal}(k_n/k) \cong \mathrm{Gal}(\ell_n/\ell) \cong \mathbb{Z}/(2^n) \text{ for all positive integers } n \in \mathbb{N}$$

---

[3] This follows, e.g., from results of Humio Ichimura and Shoichi Nakajima; see Proposition 1 and its proof in Section 3 of [IN10]

and

$$G_n := \mathrm{Gal}(\ell_n/k_n) \cong \mathbb{Z}/(2) \text{ for all indices } n \in \mathbb{N} \cup \{\infty\}.$$

Thus we have exact sequences

$$\begin{aligned} 0 \to H^1(G_n, P_{\ell_n}) &\to H^2(G_n, \mathcal{O}_{\ell_n}^\times) \to H^2(G_n, \ell_n^\times) \\ &\to H^2(G_n, P_{\ell_n}) \to H^1(G_n, \mathcal{O}_{\ell_n}^\times) \to 0, \end{aligned}$$

and we have norm maps $N_n \colon \ell_n \to k_n$ for all indices $n \in \mathbb{N} \cup \{\infty\}$. In this case, $N_n(\alpha) = \alpha\bar{\alpha} = |\alpha|^2$ is just the square modulus since the restriction of complex conjugation generates G_n for all indices $n \in \mathbb{N} \cup \{\infty\}$. Thus the images of these norms maps consist entirely of totally positive[4] elements.

Now we need a theorem. It generalizes a result of Weber's (see Sätze 6 and 25 in [Has52]) which Ferrero used in [Fer80].

Theorem 10 (I. Hughes and R. Mollin, [HM83])**.** *Let F'/F be a cyclic 2-extension of real abelian number fields. Suppose* $\mathrm{Gal}(F/\mathbb{Q})$ *has exponent n such that -1 is congruent to a power of 2 modulo n, and, if $F \neq F'$, suppose that exactly one prime ramifies in F'/F. If the class number h_F of F is odd, then every totally positive element of $\mathcal{O}_{F'}^\times$ is a square in $\mathcal{O}_{F'}^\times$.*

For all $n \in \mathbb{N}$ we apply Theorem 10 to the extension $F'/F = k_n/k$ to get

$$\mathcal{O}_{k_n}^\times \cap N_\infty(L^\times) = (\mathcal{O}_{k_n}^\times)^2 = N_n(\mathcal{O}_{\ell_n}^\times) \tag{10.1}$$

where $(\mathcal{O}_{k_n}^\times)^2$ denotes the subgroup of squares of units. In fact, taking unions shows that Equation 10.1 also holds for $n = \infty$, so

$$H^1(G, P_L) \cong \ker(\mathcal{O}_K^\times/(\mathcal{O}_K^\times)^2 \to K^\times/N_\infty(L^\times)) = (\mathcal{O}_K^\times \cap N_\infty(L^\times))/(\mathcal{O}_K^\times)^2 = 0,$$

and likewise $H^1(G_n, P_{\ell_n}) = 0$ for all $n \in \mathbb{N}$.

Hence it remains only to show that $|H^2(G, P_L)| = 2$. To do this, we first prove $|H^1(G, \mathcal{O}_L^\times)| = 2$ and then show that the surjection $H^2(G, P_L) \to H^1(G, \mathcal{O}_L^\times)$ is also an injection.

For each $n \in \mathbb{N}$, let $t_\infty(n)$ denote the number of infinite places of k_n which ramify in ℓ_n/k_n. Since k_n is totally real and ℓ_n is totally complex, $t_\infty(n) = [k_n : \mathbb{Q}]$ is just the number of real places of k_n, so Dirichlet's unit theorem gives

$$\mathcal{O}_{k_n}^\times \cong \mathbb{Z}^{t_\infty(n)+0-1} \oplus \frac{\mathbb{Z}}{(2)}$$

as abelian groups. Then using Theorem 10 again shows

$$|H^2(G_n, \mathcal{O}_{\ell_n}^\times)| = \left|\frac{\mathcal{O}_{k_n}^\times}{(\mathcal{O}_{k_n}^\times)^2}\right| = \left|\frac{\mathbb{Z}^{t_\infty(n)-1} \oplus (\mathbb{Z}/(2))}{2(\mathbb{Z}^{t_\infty(n)-1} \oplus (\mathbb{Z}/(2)))}\right| = 2^{t_\infty(n)-1+1} = 2^{t_\infty(n)}. \tag{10.2}$$

Now we state another needed result, whose proof may be found, for example, in [Gre10].

Theorem 11. *Suppose F'/F is a quadratic extension of number fields and let t_∞ denote the number of infinite places of F which ramify in F'. Then*

$$\chi(\mathrm{Gal}(F'/F), \mathcal{O}_{F'}^\times) = t_\infty - 1.$$

---

[4] Recall that an algebraic number α is called totally positive if the images of α under every embedding $\mathbb{Q}(\alpha) \hookrightarrow \mathbb{C}$ are real and positive.

We now apply Theorem 11 for the extension $F'/F = \ell_n/k_n$ to conclude that

$$|H^1(G_n, \mathcal{O}_{\ell_n}^\times)| = 2$$

for all positive integers $n \in \mathbb{N}$ since we have already shown in Equation 10.2 above that $|H^2(G_n, \mathcal{O}_{\ell_n}^\times)| = 2^{t_\infty(n)}$. On the other hand, for $n \in \mathbb{N} \cup \{\infty\}$ we have

$$U_n/V_n \cong H^1(G_n, \mathcal{O}_{\ell_n}^\times)$$

where U_n is the norm 1 units in $\mathcal{O}_{\ell_n}$ and $V_n = \{\bar{u}/u : u \in \mathcal{O}_{\ell_n}^\times\}$. We claim U_n/V_n is generated by the coset of -1 for all indices $n \in \mathbb{N} \cup \{\infty\}$. We need the following lemma which we do not prove here but is not hard to establish (see, for example, Lemma 6.15 in [Sch12]).

Lemma 12. *Let $d > 1$ be a squarefree integer. Suppose F is a number field with discriminant Δ_F such that $(d, \Delta_F) \leq 2$. Then $4\mathcal{O}_{F(\sqrt{-d})} \subseteq \mathcal{O}_F + \sqrt{-d}\mathcal{O}_F$.*

Suppose contrary to the claim that $-1 = \bar{u}/u \in V_n$ for some $u \in \mathcal{O}_{\ell_n}^\times$ and some positive integer $n \in \mathbb{N}$. Then both u and u^{-1} are in $\mathcal{O}_{\ell_n}$, so since $u = -\bar{u}$ and $u^{-1} = -\overline{u^{-1}}$ the lemma implies

$$u = i\frac{a\sqrt{d}}{4} \text{ and } u^{-1} = i\frac{b\sqrt{d}}{4}$$

for some $a, b \in \mathcal{O}_{k_n}$. Hence $abd = -4^2$, so d divides $4^2 = 2^4$ in $\mathcal{O}_{k_n}$, but that means d divides 2^4 in $\mathbb{Z}$. Therefore $d = 1$ or $d = 2$ since d is a squarefree positive integer, which contradicts our assumption that $d > 2$. Thus for each $n \in \mathbb{N}$ we know that $-1 \notin V_n$ and $|U_n/V_n| = 2$, so U_n/V_n is generated by the coset of -1. It follows that $H^1(G, \mathcal{O}_L^\times) \cong U_\infty/V_\infty$ is also generated by the coset of -1 and has order 2 since

$$U_\infty = \bigcup_{n\in\mathbb{N}} U_n = \bigcup_{n\in\mathbb{N}} \langle -1\rangle V_n = \langle -1\rangle V_\infty$$

and

$$-1 \notin \bigcup_{n\in\mathbb{N}} V_n = V_\infty.$$

To summarize, we have an exact sequence

$$0 \to H^2(G, \mathcal{O}_L^\times) \to H^2(G, L^\times) \to H^2(G, P_L) \to H^1(G, \mathcal{O}_L^\times) \to 0$$

where $|H^1(G, \mathcal{O}_L^\times)| = 2$, so to prove $|H^2(G, P_L)| = 2$ (and thus finish the proof of Theorem 8) we only need to show that the map $H^2(G, \mathcal{O}_L^\times) \to H^2(G, L^\times)$ is onto since that would imply the map $H^2(G, L^\times) \to H^2(G, P_L)$ is trivial and, consequently, that the map $H^2(G, P_L) \to H^1(G, \mathcal{O}_L^\times)$ is bijective.

We have a commutative square

$$\begin{tikzcd}
H^2(G, \mathcal{O}_L^\times) \arrow[r] \arrow[d, "\wr"] & H^2(G, L^\times) \arrow[d, "\wr"] \\
\varinjlim_n \dfrac{\mathcal{O}_{k_n}^\times}{N_n(\mathcal{O}_{\ell_n}^\times)} \arrow[r] & \varinjlim_n \dfrac{k_n^\times}{N_n(\ell_n^\times)}
\end{tikzcd}$$

where the horizontal maps are the natural maps and the vertical maps are isomorphisms. Pick some $n \in \mathbb{N}$ and $x_n \in k_n^\times$. Then the commutative square above implies that proving the surjectivity of $H^2(G,\mathcal{O}_L^\times) \to H^2(G,L^\times)$ amounts to showing there is an $m \in \mathbb{N}$ and a $y_m \in \mathcal{O}_{k_m}^\times$ such that we have an equality of cosets $x_n N_j(\ell_j^\times) = y_m N_j(\ell_j^\times)$ for some integer $j \geq \max\{n,m\}$. In fact, we show that we can take $j = m+1$.

For the moment, let m be an arbitrary positive integer. Define S_m to be the set of places of k_m which do not split in ℓ_m. We have a commutative diagram

(12.1)

$$\begin{tikzcd}
\dfrac{\mathcal{O}_{k_{m+1}}^\times}{N_{m+1}(\mathcal{O}_{\ell_{m+1}}^\times)} \arrow[r, "\beta_{m+1}"] & \dfrac{k_{m+1}^\times}{N_{m+1}(\ell_{m+1}^\times)} \arrow[r, "\sim"] & \bigoplus_{v' \in S_{m+1}}' \mathrm{Br}(\ell_{m+1,w'}/k_{m+1,v'}) \\
\dfrac{\mathcal{O}_{k_m}^\times}{N_m(\mathcal{O}_{\ell_m}^\times)} \arrow[u, "\alpha_m"] \arrow[r, "\beta_m"] & \dfrac{k_m^\times}{N_m(\ell_m^\times)} \arrow[u, "\gamma_m"] \arrow[r, "\sim"] & \bigoplus_{v \in S_m}' \mathrm{Br}(\ell_{m,w}/k_{m,v}) \arrow[u, "\delta_m"]
\end{tikzcd}$$

where α,β,δ are the natural maps and the restricted direct sum $\oplus'$ on relative local Brauer groups contains only those tuples which lie in the kernel of the natural map to $\mathbb{Q}/\mathbb{Z}$ given by the sum of local invariants.[5] We have already noted that β_m, β_{m+1} are injective, and α_m is injective for the same reason, i.e., Theorem 10 implies

$$\mathcal{O}_{k_m}^\times \cap N_{m+1}(\mathcal{O}_{\ell_{m+1}}) = (\mathcal{O}_{k_m}^\times)^2 = N_m(\mathcal{O}_{\ell_m}^\times).$$

Thus $\beta_{m+1} \circ \alpha_m$ is injective, so commutativity of the left side of Diagram 12.1 and Equation 10.2 imply

$$|\mathrm{im}(\gamma_m \circ \beta_m)| = |\mathrm{im}(\beta_{m+1} \circ \alpha_m)| = |\mathcal{O}_{k_m}^\times / N_m(\mathcal{O}_{\ell_m}^\times)| = 2^{t_\infty(m)}. \tag{12.2}$$

Let H_m be the subgroup consisting of those tuples in $\bigoplus_{v\in S_m}' \mathrm{Br}(\ell_{m,w}/k_{m,v})$ which are trivial on the components corresponding to finite places $v \in S_m$ which split in k_{m+1}. We can show $|\delta_m(H_m)| = 2^{t_\infty(m)}$ using the fact that only the infinite components of H_m will not be killed by δ_m.

Now we specify m. Finite places are finitely split in the cyclotomic $\mathbb{Z}_p$-extension of a number field, so there is a positive integer m for which

$$x_n N_m(\ell_m^\times) \in \tilde{H}_m \tag{12.3}$$

where $\tilde{H}_m$ denotes the subgroup of $k_m^\times / N_m(\ell_m^\times)$ which maps isomorphically onto H_m in the above commutative diagram 12.1. We have $|\gamma_m(\tilde{H}_m)| = |\delta_m(H_m)| = 2^{t_\infty(m)}$ and we will show that $\mathrm{im}(\gamma_m \circ \beta_m) \subseteq \gamma_m(\tilde{H}_m)$, so Equation 12.2 implies via the pigeonhole principle that

$$\gamma_m(\tilde{H}_m) = \mathrm{im}(\gamma_m \circ \beta_m). \tag{12.4}$$

To see $\mathrm{im}(\gamma_m \circ \beta_m) \subseteq \gamma_m(\tilde{H}_m)$, we first note that the study of Brauer groups for local fields shows $\mathrm{im}(\beta_m)$ maps injectively into the subgroup of $\bigoplus_{v\in S_m}' \mathrm{Br}(\ell_{m,w}/k_{m,v})$ consisting of tuples which are trivial on all components other than those corresponding to infinite places and to finite places which ramify in ℓ_m/k_m. Increasing m if necessary, we may assume that the finite places $v \in S_m$ which ramify in ℓ_m/k_m do not split in k_{m+1}. This means that $\operatorname{im}(\beta_m) \subseteq \tilde{H}_m$, and hence $\operatorname{im}(\gamma_m \circ \beta_m) \subseteq \gamma_m(\tilde{H}_m)$, as needed. Finally, Equations 12.3 and 12.4 give

---

[5] Note that $v \in S_m$ uniquely specifies a place w of ℓ_m which lies above v. Also, if $v \notin S_m$ is a place of k_m, then $\mathrm{Br}(\ell_{m,w}/k_{m,v})$ is trivial for any place w of ℓ_m lying over v.

$$x_n N_{m+1}(\ell_{m+1}^\times) = \gamma_m(x_n N_m(\ell_m^\times)) = \gamma_m(\beta_m(y_m N_m(\mathcal{O}_{\ell_m}^\times))) = y_m N_{m+1}(\ell_{m+1}^\times)$$

for some $y_m \in \mathcal{O}_{k_m}^\times$, which finishes the proof. □

References

[Die40] Fritz-Erdmann Diederichsen, *Über die Ausreduktion ganzzahliger Gruppendarstellungen bei arithmetischer Äquivalenz*, Abh. Math. Sem. Hansischen Univ. **13** (1940), 357–412.

[Fer80] Bruce Ferrero, *The cyclotomic $\mathbb{Z}_2$-extension of imaginary quadratic fields*, Amer. J. Math. **102** (1980), no. 3, 447–459.

[FW79] Bruce Ferrero and Lawrence C. Washington, *The Iwasawa invariant μ_p vanishes for abelian number fields*, Ann. of Math. (2) **109** (1979), no. 2, 377–395.

[Gre10] Ralph Greenberg, *Topics in Iwasawa theory*, math.washington.edu/, 2010.

[Has52] Helmut Hasse, *Über die klassenzahl abelscher zahlkörper*, Akademie Verlag, Berlin, 1952.

[HM83] I. Hughes and R. Mollin, *Totally positive units and squares*, Proceedings of the American Mathematical Society **87** (1983), no. 4, 613–616 (English).

[IN10] Humio Ichimura and Shoichi Nakajima, *On the 2-part of the ideal class group of the cyclotomic $\mathbb{Z}_p$-extension over the rationals*, Abhandlungen aus dem Mathematischen Seminar der Universität Hamburg **80** (2010), 175–182.

[Iwa59] Kenkichi Iwasawa, *On Γ-extensions of algebraic number fields*, Bull. Amer. Math. Soc. **65** (1959), 183–226.

[Iwa65] Kenkichi Iwasawa, *Analogies between number fields and function fields*, Annual science conference proceedings - Yeshiva University, Belfer Graduate School of Science **2** (1965), 203–208.

[Iwa73a] Kenkichi Iwasawa, *On $\mathbf{Z}_l$-extensions of algebraic number fields*, Ann. of Math. (2) **98** (1973), 246–326.

[Iwa73b] ———, *On the μ-invariants of Z_1-extensions*, Number theory, algebraic geometry and commutative algebra, in honor of Yasuo Akizuki, Kinokuniya, Tokyo, 1973, pp. 1–11.

[Iwa81] Kenkichi Iwasawa, *Riemann-Hurwitz formula and p-adic Galois representations for number fields*, Tohoku Math. J. (2) **33** (1981), no. 2, 263–288.

[Kid79] Yûji Kida, *On cyclotomic $\mathbb{Z}_2$-extensions of imaginary quadratic fields*, Tohoku Math. J. **31** (1979), 91–96.

[Sch12] Jordan Schettler, *The Change in Lambda Invariants for Cyclic p-Extensions of $\mathbb{Z}_p$-Fields*, Ph.D. thesis, The University of Arizona, 2012.